

LIC Insert Preparation Manual

Version Number : 1

Start Production Date: 09.03.02

Author: Hisao Yokota

Edited by:

Reviewed by:

Summary: This protocol describes the standard procedure for preparing LIC Insert Stocks. Qiagen Kit purified Target PCR, 1 ug, is digested with T4 DNA polymerase exonuclease digestion in the presence of dTTP yields LIC sticky ends

Materials/Reagents/Equipment	Vendor	Stock Number
Disposables		
PCR tubes	Marsh, Perkin-Elmer	
Reagents		
T4 DNA polymerase	New England Biolabs	
dTTP mixture, 100 mM, -20C	Pharmacia, Promega	
Water, double deionized, autoclaved, RT		
Agarose	Amresco,	
TAE Buffer	BSGC	
I Kb Plus DNA ladder	Invitrogen	
Equipment		
PCR Machine 2400 or 9600	Perkin-Elmer	
Pipetman		
Agarose Gel Electrophoresis	Invitrogen	

Procedure for 20 ul Reaction

Label PCR tube – **only on upper sides** (otherwise ink stains temp block).

Setup PCR machine to run 37° C for 30 min, then 70° C for 20 min, then 4° C termination.

T4 DNA Polymerase Treatment: Add 0.5 ul of dTTP [100 mM], **0.5 ul of BSA** [10 mg/ml], mix and then add 0.5 ul [20U] T4 DNA Pol. Incubate in PCR machine cycle above.

Usage: use 1.0 ul [20 ng Insert DNA] for each LIC reaction.

LIC Insert Preparation Datasheet

Name		Date	

No.	Sample		